



## Seaweed biomass of the Philippines: Sustainable feedstock for biogas production



Gian Powell B. Marquez<sup>a,\*</sup>, Wilfred John E. Santiañez<sup>c</sup>, Gavino C. Trono Jr.<sup>c</sup>,  
Marco Nemesio E. Montaño<sup>c</sup>, Hiroshi Araki<sup>b</sup>, Hisae Takeuchi<sup>b</sup>, Tatsuya Hasegawa<sup>a,b</sup>

<sup>a</sup> Department of Aerospace Engineering, Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan

<sup>b</sup> EcoTopia Science Institute, Nagoya University, Nagoya 464-8603, Japan

<sup>c</sup> The Marine Science Institute, University of the Philippines, Diliman, Quezon City 1101, Philippines

### ARTICLE INFO

#### Article history:

Received 20 March 2014

Received in revised form

19 June 2014

Accepted 6 July 2014

Available online 26 July 2014

#### Keywords:

Anaerobic digestion

Biogas

Methane

Methanogens

Philippines

Seaweed

### ABSTRACT

Archipelagic and island countries are among the most affected by the worsening climate change. To mitigate global warming, the Philippines mandated the use of biofuel through the Biofuel Act of 2006. However, coping with the increasing biomass demand for biofuel production is challenging because of the country's limited freshwater and land resources. On the other hand, the Philippines has rich but untapped marine biomass resources. Hence, the potential of seaweed resources as the most accessible energy source for coastal and island communities of the Philippines was explored for methane fermentation. The seaweed species, *Sargassum* spp., *Turbinaria* spp., *Hydroclathrus* spp., *Caulerpa* spp., and *Ulva* spp., which can allow sustainable biomass supply either through managed harvesting of natural stocks or through cultivation, have been identified. Information on the geographical distribution and seasonal productivity of these species were laid out. The proximate and elemental components of the biomass were presented for viability assessment as feedstock. Challenges on biogas technology in the Philippines and anaerobic digestion of seaweed biomass were also evaluated. Research on the development of culture technologies for the *Sargassum* spp., *Turbinaria* spp., *Hydroclathrus* spp. and *Ulva* spp. is recommended to avoid the adverse impacts of harvesting on natural stocks. Further studies on anaerobic digestion conditions of the selected seaweeds that are suitable for utilization by island communities should also be conducted.

© 2014 Elsevier Ltd. All rights reserved.

### Contents

1.	Introduction	1057
2.	Seaweed biomass: prospects for sustainable feedstock	1058
2.1.	The genus <i>Sargassum</i>	1058
2.1.1.	Biomass distribution and production	1058
2.2.	The genus <i>Turbinaria</i>	1060
2.2.1.	Biomass distribution and production	1060
2.3.	The genus <i>Hydroclathrus</i>	1060
2.3.1.	Biomass distribution and production	1060
2.4.	The genus <i>Caulerpa</i>	1060
2.4.1.	Biomass distribution and production	1060
2.5.	The genus <i>Ulva</i>	1060
2.5.1.	Biomass distribution and production	1060
3.	Biogas production process	1060
3.1.	Anaerobic digestion process	1060
3.1.1.	Hydrolysis	1061
3.1.2.	Acidogenesis	1061

\* Correspondence to: Room 524, Hasegawa's Laboratory, EcoTopia Science Institute, Department of Aerospace Engineering, Graduate School of Engineering, Nagoya University, Nagoya 464-8603, Japan. Tel.: +81 52 789 5487; fax: +81 52 789 4506.

E-mail address: [g.p.b.marquez@yuki.esi.nagoya-u.ac.jp](mailto:g.p.b.marquez@yuki.esi.nagoya-u.ac.jp) (G.P.B. Marquez).

3.1.3.	Acetogenesis.....	1061
3.1.4.	Methanogenesis.....	1061
4.	Biomethane potential of seaweed biomass.....	1062
4.1.	Properties of seaweed biomass.....	1062
4.1.1.	Elemental components.....	1062
4.1.2.	Proximate compositions.....	1063
4.2.	Methane fermentation of seaweed biomass.....	1063
5.	Engaging coastal communities on green energy.....	1064
6.	Conclusions.....	1064
	Acknowledgment.....	1065
	References.....	1065

## 1. Introduction

Oil crisis in the 1970s drove the development of biogas technology [1] using several species of macroalgae [2,3] as one of the renewable energy solutions. Interest in renewable energy waned when oil prices stabilized because of the discovery of new oil fields. Recently, the interests in marine biomass utilization for biofuel production reemerged due to several reasons. These are (1) mitigation of climate change by lowering CO<sub>2</sub> emissions, and maintaining the closed carbon cycle [4], (2) energy security by decreasing the reliance on petroleum fuels, and its erratic price fluctuation [5], (3) food security by avoiding the indirect and direct competition on terrestrial food crops utilization for biofuel production, and preventing high food prices [6], (4) better feedstock compared to terrestrial crops by having faster growth rate [7], and (5) environmental conservation and preservation by preventing forest conversion to biofuel crop farms [8].

The Philippines is among the most impacted countries of global warming as shown on its global risk index rank – from third place in 2009 [9] to second place in 2012 [10]. To mitigate climate change and its inevitable risks and impacts, the Philippine government institutionalized the Biofuel Act of 2006 (Republic Act 9367), which mandates the blending of gasoline with bioethanol from 10% in 2012 to 20% in 2025 and of diesel with biodiesel from 2% in 2012 to 10% in 2025 [11]. The Energy Regulatory Commission also approved a feed-in-tariff of ₱6.63/kWh (~US\$0.15) for electricity generated from biomass renewable energy last July 2012 [12]. The support from the different sectors in rapidly developing the bioethanol and biodiesel industries helps in achieving the clean energy goals of the Philippines. Nevertheless, whether or not these efforts are sustainable in the long term is still uncertain.

Biodiesel and bioethanol are liquid biofuels that are preferred commercially because these are easy to transport and can be readily mixed with petroleum gasoline and diesel [13]. Coconut oil supplies most of the biodiesel demand in the Philippines, but other feedstock such as *Jatropha* and palm oil are also explored [14]. On the other hand, bioethanol production heavily relies on sugarcane, but it can only supply 79% of the country's need [14]. The demand for bioethanol production may continue to increase with the encouragement of the Philippine government on biofuel use. However, the sustainability of these mechanisms remains uncertain. The conversion of arable and traditional food crop-producing farms to biofuel feedstock production may negatively affect food security, considering the steady growth of the Philippine population. Decrease in food crop production may also increase the price of food, reducing the accessibility of the masses. Moreover, large-scale mono-cropping for biofuel feedstock may have negative environmental effects such as biodiversity loss, which is in conflict with various national policies and mechanisms on environmental conservation and protection. These factors can

pose a problem in achieving the country's goal on biofuel sustainability.

Among the biomass-utilizing biofuel technologies (e.g., bioethanol fermentation and biodiesel production), biogas production is the most efficient in terms of net energy gain [15] because anaerobic digestion utilizes all the degradable components (carbohydrate, protein and lipid) of the feedstock [16] to produce biogas (~60% CH<sub>4</sub>, ~40% CO<sub>2</sub> and other trace gases). Biogas can also be upgraded to increase its methane content up to the same level as that of natural gas, giving the same performance when used as fuel in internal combustion engines or as household cooking gas. The limited land resources of the Philippines for biomass cultivation made biogas technology important to its long term green energy goal because of the wider range of crops suitable for biogas production, as compared to biodiesel and bioethanol.

The archipelagic nature of the Philippines isolates many islands and coastal communities, resulting in the limited access of these areas to a centralized power source. Despite the government's increased efforts to provide electricity for its people, many distant and poor communities still have insufficient, if not complete absence of, electricity supply. The use of household biogas digesters for the production of biogas as an alternative and climate-friendly fuel source may be suitable and practical, especially for poor families residing in these many and widespread isolated communities.

The Philippines' coastline is 36,289 km long – the second longest in Asia and the fourth in the world [17]. However, the country's abundant marine resources, especially the often conspicuous seaweed resources, remain relatively unexplored and underdeveloped for the sustainable production of biofuel feedstock. Seaweeds have long been part of the lives of coastal communities in the Philippines. Seaweeds can be categorized on the basis of its use, namely, as food, source of phyco-colloids, and source of other natural products used in various applications [18]. Out of the 350 species (820 species listed for the country) that have known economic values, less than 5% are commercially important. Among these commercially important species, only the phycocolloid-producing species, the agarophytes (*Gracilaria* spp. and *Gelidiella acerosa*) and the carrageenophytes (*Eucheuma* spp. and *Kappaphycus* spp.), gained considerable attention. The potential of other seaweed resources of the country remains untapped, and the majority remains underdeveloped or undeveloped [18].

Although seaweed resource utilization in the Philippines is mostly focused on natural product extraction, the increasing interest in the utilization of these biomasses for biofuel, specifically biogas production, instigated this review. Hence, this study aimed to determine the different seaweed species in the Philippines that can be sustainably utilized as biogas feedstock for commercial or household use. The biomass productivity, as

well as composition and degradability of the seaweeds are also discussed. Research on the anaerobic digestion of marine biomass suitable for utilization by coastal and island folks is presented.

## 2. Seaweed biomass: prospects for sustainable feedstock

The criteria for the selection of species for biofuel production generally include the following: (1) the species must be available in large quantities in many areas of the country, that is, it should sustainably provide for the requirements of production; (2) the species should be available year round or in most part of the year; and, (3) sustainable cropping (or cultivation) can be conducted without adverse effects on the environment. Several species satisfy at least one, if not all, of these criteria. These species are the brown seaweeds *Sargassum* spp., *Turbinaria* spp., and *Hydroclathrus* spp. and the green seaweeds *Caulerpa* spp. and *Ulva* spp. (Fig. 1). Although members of the genus *Eucheuma*, *Kappaphycus*, *Gracilaria* and *Gelidiella* may also satisfy these criteria, these seaweeds are more economically valuable as feedstock for the production of natural products than for biofuel production.

### 2.1. The genus *Sargassum*


The members of the genus *Sargassum* C. Agardh (Phylum Ochrophyta, Class Phaeophyceae, Order Fucales, and Family Sargassaceae) are widely distributed and common in the Indo-Pacific region. Currently, this genus has 335 taxonomically accepted species [20]. At least 29 of the more than 90 species

reported in the Philippines have been validated. Members of this genus are conspicuous and form extensive beds along the rocky shores of the country.

In addition to its high ecological value, *Sargassum* is well known for its economically important natural products such as alginates, fucoidan, fucoxanthin, and lutein. The ethnobotanical uses of *Sargassum* biomass by coastal populations across the country include its use as a cover for the prevention of desiccation and maintenance of freshness of fishery products, food, plant fertilizer and flower inducer, insect repellent, animal feed in agriculture, and therapeutic drink, among others [21]. *Sargassum* biomass also has great potential as feedstock for biofuel [22].

#### 2.1.1. Biomass distribution and production

*Sargassum* biomass production in the Philippines, as in any part of the world, coincides with the growth and development of the species (i.e., period of regeneration or recruitment, fast growth phase, reproduction, senescence, and die-back) [23–32]. According to various studies conducted in the country, *Sargassum* biomass production is low during recruitment, but increases during the fast growth phase. Biomass production peaks during reproduction, and consequently decreases at the onset of senescence and die-back. During the latter stage, much of the biomass produced are uprooted and washed ashore. These dislodged seaweeds, together with sea grasses (collectively known as sea wrack), can be collected and used for biogas production as demonstrated by Marquez et al. [33]. Moreover, in the absence of technology for the mariculture of *Sargassum*, a cropping management scheme for existing *Sargassum* beds was developed by Trono and Tolentino [34] to maximize the utilization and conservation of the available harvestable biomass from natural beds. Intertidal populations can



**Fig. 1.** The seaweed resources of the Philippines ((a) *Sargassum* bed; (b) *Caulerpa lentillifera*; (c) *Turbinaria conoides*; (d) *Hydroclathrus clathratus*; and (e) *Ulva reticulata*) with high potential as feedstock for biogas production. Photo courtesy of (a) WJE Santiañez and (b–e) Trono [19].



be harvested once a year, and subtidal populations can be harvested at least twice. Harvests can be scheduled before fertility peak or when approximately 50% of the populations are fertile. This strategy allows the species to regenerate and proceed to the regular reproductive phase. Moreover, pruning must be conducted near the base, ensuring that both the holdfast and primary laterals are left to avoid interference with the regeneration process [23].

Harvest must also be conducted in strips or randomly to ensure enough fertile materials for recruitment [34].

Studies on *Sargassum* biomass production in the Philippines are summarized in Table 1. Many of the *Sargassum* beds produce high biomass during the last quarter of the year (November to December). Given these information, insights on harnessing the potential of *Sargassum* biomass for biogas production can be derived.

**Table 1**

Biomass production of *Sargassum* species in the Philippines.

Species	Highest biomass (kg wet weight/m <sup>2</sup> )	Period	Locality	Ref.
<i>S. binderi</i>	3.36		Mactan Is., Cebu	[35]
<i>S. crassifolium</i>	0.667		Taklong Is., Guimaras	[36]
	2.55	August	Maydolong, Samar	[37]
	13.19	January/February	Calarian, Zamboanga	[38]
	2.98	January/February	Bolinao, Pangasinan	
<i>S. cristaefolium</i>	0.647	December	Bolinao, Pangasinan	[39]
	0.430	November	Bolinao, Pangasinan	
	0.242	September	Ilocos Sur	[40]
	0.111 <sup>a</sup>	February	Negros Island	[26]
<i>S. feldmanii</i>	0.723 <sup>a</sup>	March	Negros Island	
<i>S. hemiphyllum</i>	0.963	March	Maydolong, Samar	[37]
<i>S. ilicifolium</i>	0.124 <sup>a</sup>	February	Negros Island	[26]
<i>S. oligocystum</i>	15.60	January/February	Calarian, Zamboanga	[38]
	2.38	October	Maydolong, Samar	[37]
	0.282	November	Bolinao, Pangasinan	[39]
	1.60	December	Maydolong, Samar	[37]
<i>S. polycystum</i>	0.011 <sup>a</sup>	May	Negros Island	[26]
	0.447	December	Bolinao, Pangasinan	[39]
	3.07	July	Liloan, Cebu	[41]
	11.09	January/February	Quezon, Palawan	[38]
<i>S. siliquosum</i>	12.08	January/February	Aborlan, Palawan	
	3.51	December	Maydolong, Samar	[37]
	1.24	August	Pasacao, Camarines Sur	[42]
	4.30		Danajon Reef, Bohol	[43]
<i>S. siliquosum</i>	6.61	December	Liloan, Cebu	[41]
	11.36	January/February	Aborlan, Palawan	[35]
	2.69		Mactan Is., Cebu	
	4.89	January	Maydolong, Samar	[37]
<i>Sargassum</i> spp.	0.478 <sup>a</sup>	August	Calatagan, Batangas	[44]
	1.08		Calarian, Zamboanga	[35]
	7.16		Aborlan, Palawan	
	4.70		Tadtaran, Quezon	
	4.14		Cidanao, Quezon	
	4.25		Bolinao-Anda, Pangasinan	[43]
	3.20		Currimao, Ilocos Norte	
	8.40		Burgos, Ilocos Norte	
	3.81		Nasugbu Bay, Batangas	
	3.00		Matabungkay, Batangas	
	0.335 <sup>a</sup> (intertidal)	March	Currimao, Ilocos Norte	[45]
	0.600 <sup>a</sup> (subtidal)	July		

<sup>a</sup> Indicates dry weight values.

**Table 2**

Biomass production of the seaweed genera *Caulerpa*, *Ulva*, *Hydroclathrus*, and *Turbinaria* in the Philippines.

Species	Highest biomass (kg wet weight/m <sup>2</sup> )	Period	Locality	Ref
<b>Chlorophyta</b>				
<i>C. lentillifera</i>	1.53	April–May (pond culture)	Calatagan, Batangas	[51]
<i>U. lactuca</i>	2.60	March	Mactan Is., Cebu	[52]
<i>U. reticulata</i>	0.150	March	Mactan Is., Cebu	
<b>Phaeophyceae</b>				
<i>H. clathratus</i>	0.204	April	Taklong Is., Guimaras	[36]
	0.370	February	Ilocos Sur	[40]
	0.017	April	Pasacao, Camarines Sur	[42]
<i>T. conoides</i>	0.421	October	Taklong Is., Guimaras	[36]
<i>T. ornata</i>	0.291	September	Pasacao, Camarines Sur	[42]
	0.278	October	Taklong Is., Guimaras	[36]
<i>T. luzonensis</i>	0.128	October	Pasacao, Camarines Sur	[42]
	0.116	April	Taklong Is., Guimaras	[36]
<i>Turbinaria</i> sp.	1.34		Mactan Is., Cebu	[35]

## 2.2. The genus *Turbinaria*

The members of the genus *Turbinaria* Lamoroux (Phylum Ochrophyta, Class Phaeophyceae, Order Fucales and Family Sargassaceae [20]) are generally found attached on rocks or sandy-coralline substrates in reef areas with moderate to strong water movement [46]. This genus has at least 22 species (including forms and varieties) in the world [20], most of which are widely distributed in the tropics. At least 10 species (excluding varieties and forms) can be found in the Philippines [47].

### 2.2.1. Biomass distribution and production

*Turbinaria conoides*, *T. ornata*, *T. decurrens* and *T. luzonensis* can be found in many coastal areas in the country. All these species, except the latter, are common in the intertidal and subtidal areas of the coasts. Members of the genus *Turbinaria*, especially *T. conoides*, can form thick, huge colonies in subtidal areas [19]. *Turbinaria* beds are among the most conspicuous marine flora in the country, next to *Sargassum* [48].

Studies on the *Turbinaria* species in the country are very limited. In Pasacao, Camarines Sur, the biomass produced by *T. ornata* and *T. luzonensis* peaks on September (0.45 kg/m<sup>2</sup>) and October (0.12 kg/m<sup>2</sup>), respectively [42]. *Turbinaria* found in Mactan Island, Cebu has a fresh biomass of 1.34 kg/m<sup>2</sup> [35]. Other pertinent information on the seasonal distribution of the different species across the country can be found in Trono [19].

## 2.3. The genus *Hydroclathrus*

The members of the genus *Hydroclathrus* Bory de Saint-Vincent (Phylum Ochrophyta, Class Phaeophyceae, Order Ectocarpales and Family Scytosiphonaceae [20]) are common in tropical waters but highly seasonal. Peak in abundance occurs during summer and is marked by the formation of dense mats, covering most of the substrate and other benthic organisms [19,46]. Biomass decreases at the onset of the rainy season, leading to complete absence [19]. *Hydroclathrus* has four known species in the world [20], two of which can be found in the Philippines, namely *Hydroclathrus clathratus* and *H. tenuis* [19].

### 2.3.1. Biomass distribution and production

*Hydroclathrus clathratus* and *H. tenuis*, are commonly found on reef flats, sometimes attached to rocks or interspersed among sea grasses. Among these two species, *H. clathratus* is more common and abundant, but both are seasonal. Both species are abundant during the period of warmer waters (summer), in the months of March to May or June [19,49]. San Diego-McGlone et al. [50] noted the bloom of *H. clathratus* in Puerto Galera, linking them to nutrient mediated stress (nutrient enrichment). Despite its extreme seasonality, much of the biomass produced by *Hydroclathrus* spp. is underutilized and hence holds great potential for biogas production. However, the scarcity of information on the field biology and ecology of this seaweed may hamper its optimum utilization. The available information on the biomass production capacity of both species in the country is listed in Table 2. Trono [19] summarized the information on the country-wide seasonality of *H. clathratus* and *H. tenuis*.

## 2.4. The genus *Caulerpa*

The members of the genus *Caulerpa* Lamoroux (Phylum Chlorophyta, Class Bryopsidophyceae, Order Bryopsidales and Family Caulerpacae [20]) are widespread, occurring in tropical to sub-tropical waters [53]. The species are known to thrive in calm, sheltered or moderately exposed areas generally with sandy

substratum but can also be found growing on muddy, sandy-coralline, and rocky substrates [46]. At least 24 species of *Caulerpa* can be found in the Philippines, excluding the different varieties of each species [47]. Most species are commercially consumed as human food, especially as seaweed salad [19,54].

### 2.4.1. Biomass distribution and production

Most species (*Caulerpa cupressoides*, *C. lentillifera*, *C. peltata*, and *C. racemosa*) form colonies or wide mats on sandy, muddy or rocky substrates. Among all the species, *C. lentillifera* is the most favored species for consumption; hence, this species is commercially farmed in ponds and lagoons [19]. The farming technology for the mass culture of *C. lentillifera* that was developed in the 1980s [51] provided greater prospect in the utilization and sustainable production of the species as feedstock for biofuel production. Maximum annual production of *C. lentillifera* grown in a 10,000 m<sup>2</sup> farm at a 0.10 kg/m<sup>2</sup> initial seeding may reach up to 1.53 kg/m<sup>2</sup> wet weight [51]. The temporal and spatial distribution of various *Caulerpa* species was documented by Trono [19,54].

## 2.5. The genus *Ulva*

The members of the genus *Ulva* Linnaeus (Phylum Chlorophyta, Class Ulvophyceae, Order Ulvales, and Family Ulvaceae [20]) are common in the tropics, but can also be found in the subtropical waters of China. The species are often found in shallow intertidal areas that are intermittently exposed to air during low tide. The species can also be found as epiphytes of other large marine algae such as *Sargassum* or on sea grass blades [46]. A total of 98 species can be found in the world [20], 12 of which are found in the Philippines [47], excluding infraspecific names.

### 2.5.1. Biomass distribution and production

Most species are characteristic flora of eutrophic waters and considered as pollution indicators [52]. Although it can be found all throughout the year, *Ulva* spp. are more abundant from November to April. However, little is known on the actual amount of biomass produced in the different localities across the country. The only report that quantified the biomass of *Ulva* spp. is that of Largo et al. [52] in Mactan, Cebu, where the species form extensive beds (i.e., "green tide") during summer (March) and has fresh biomass reaching up to 2.6 kg/m<sup>2</sup> (Table 2).

## 3. Biogas production process

Most studies on biofuel production using marine biomass focused on microalgae for biodiesel [55,56] and macroalgae for bioethanol production [57,58]. Biodiesel and bioethanol are liquid biofuels that are commercially preferred because it can readily and easily utilized by different vehicle and industrial engines for power generation. Although biogas can be directly used with gas stoves in households as a substitute for liquefied petroleum gas, it needs further upgrading for commercial engine usage. Therefore, biogas is preferred as a renewable source of energy in rural communities [59,60]. Household biogas digesters can be easily managed and operated without the need for complicated training for the users. A wider range of biomass can also be used as feedstock. Nonetheless, the anaerobic digestion process for the production of biogas involves complex interactions between diverse microorganisms and is discussed below.

### 3.1. Anaerobic digestion process

The anaerobic digestion process has four phases, namely, hydrolysis, acidogenesis, acetogenesis, and methanogenesis




Fig. 2. The four phases of anaerobic digestion in biogas production showing the different organisms involved [60–64].

(Fig. 2) [61–64]. The hydrolysis, acidogenesis, and acetogenesis phases are collectively called the acidification step because of the drastic decrease in pH resulting from the fast occurrence of these phases compared with that of the methanogenesis phase [65]. The faster growth rate and lower physico-chemical sensitivity of the microorganisms in the acidification step commonly cause instability to the anaerobic digestion process.

### 3.1.1. Hydrolysis

In hydrolysis, strictly or facultatively anaerobic bacteria secrete extracellular enzymes that break down polymers of organic matter such as protein, polysaccharides, and lipids into their respective monomer and oligomer states. The anaerobic digestion of different biomasses completely depends on the capability of the microorganisms present as microflora or as microbial seed to produce appropriate hydrolytic enzymes for specific polymers. Amylolytic, proteolytic, cellulolytic, lipolytic, or phycocolloid-hydrolyzing bacteria can effectively hydrolyze starchy crops, protein-rich waste products, cellulosic plants, fatty fruits, or phycocolloid-abundant seaweeds biomass, respectively.

Lignocellulosic polymers are difficult to degrade; hence, pre-treatment is needed to break down their complex structures and make cellulose accessible. Lignin cannot be degraded anaerobically because lignin-depolymerizing enzymes require oxygen. Before proceeding to acidogenesis, cellulose and hemicellulose, except lignin, are first digested by hydrolytic bacteria. Some microorganisms that can be found in an anaerobic digester include *Clostridia* species; this microorganism has cellulosome which is a multi-enzyme complex consisting of endo-1, 4- $\beta$ -glucanase, exo-1, 4- $\beta$ -glucanase, and  $\beta$ -galactosidase – that facilitates adhesion and complete hydrolysis of cellulose to glucose [66]. Hemicellulose such as xylan can be hydrolyzed into pentose sugar and xylose by some microbes such as *Caldicellulosiruptor saccharolyticus* [67].

### 3.1.2. Acidogenesis

Acidogenesis is the second phase in the anaerobic digestion process. The monomer and oligomer end products from the hydrolysis are instantly consumed in this phase by fermentative acidogens (strictly or facultatively anaerobic). Three major end-product substrates are produced in this phase and are grouped into (1) acetate, (2) hydrogen/carbon dioxide ( $H_2/CO_2$ ), and (3) non-acetate short chain fatty acids, alcohols, and methylated

substrates (NAM). Monomeric glucose from hydrolysis is converted into acetate, propionate, butyrate, ethanol [68], or methane with co-production of carbonic acid [62]. In the presence of xylose,  $H_2/CO_2$  and acetate are produced by *Caldicellulosiruptor saccharolyticus* [67], among others, which simultaneously forms a syntrophic relationship with hydrogenotrophic methanogens [69].

### 3.1.3. Acetogenesis

The third phase, acetogenesis, is where the  $H_2/CO_2$  and NAM substrates from acidogenesis are utilized by homoacetogens and syntrophic acetogens, respectively, for acetate conversion. Homoacetogens reduce  $CO_2$  into acetate using  $H_2$  as electron donor. In some cases, when population of aceticlastic methanogen, which consumes acetate in methanation step, is low, backward utilization of acetate substrate to produce  $H_2$  and  $CO_2$  is favored by homoacetogens in the presence of hydrogenotrophic methanogens. The latter simultaneously consume the  $H_2/CO_2$  produced [64,70].

Moreover, the consumption of short chain fatty acids for acetate,  $CO_2$ , and  $H_2$  production by syntrophic acetogens require an association with hydrogenotrophic methanogens. This syntrophic association allows immediate utilization of  $H_2$  by hydrogenotrophic methanogen, lowering the  $H_2$  partial pressure ( $< 10$  Pa). The simultaneous consumption of  $H_2$  permits the endergonic conversion of acetate and  $H_2/CO_2$  by releasing the electron from NADH as molecular hydrogen, inducing forward reaction in this pathway [70].

Some acetogens such as *Syntrophomonas wolfei* and *Syntrophobacter wolnii*, which consume fatty acids and propionate, respectively, are in syntrophic relation with hydrogenotrophic methanogens [62,71,72]. On the other hand, propionate and alcohols are used by *Pelotomaculum* to produce acetate and  $H_2/CO_2$  for the methanogenesis substrate [73]. Syntrophic acetogens such as *Acetobacterium woodii* [74], *Thermotoga lettingae* strain TMO [75], *Thermacetogenium phaeum* strain PB [76], and *Clostridium ultunense* strain B [77] are also associated with hydrogenotrophic methanogens [63].

### 3.1.4. Methanogenesis

There are three types of methanogens – (1) methylotrophic, (2) hydrogenotrophic and (3) aceticlastic methanogens – involved in the methanogenesis phase, which are classified based on the substrate they utilize. Methylotrophic methanogens consume the methylated substrates from the NAM that is produced during acidogenesis phase. Hydrogenotrophic methanogens utilize the  $H_2/CO_2$  substrates that are the end-products of acidogens and homoacetogens and co-products of syntrophic acetogens. Aceticlastic methanogens use acetate substrates that are the end products in acidogenesis and homoacetogenesis and co-products in syntrophic acetogenesis [61,62,64,78].

Methanogens have six taxonomic orders (i.e., Methanococcales, Methanomicrobiales, Methanobacteriales, Methanocellales, Methanopyrales, and Methanosarcinales) [79]. Four of these orders are commonly found in anaerobic digesters [80], namely, the hydrogenotrophic Methanococcales and Methanomicrobiales, which mostly use  $CO_2$  and either formate or  $H_2$  as electron donors; the hydrogenotrophic Methanobacteriales, which use mostly  $CO_2$ , and  $H_2$  as an electron donor, but some use  $CO_2$ , and both  $H_2$  and formate; and Methanosarcinales, which use acetate (aceticlastic), methanol and methylamines (methylotrophic), and  $H_2/CO_2$  (hydrogenotrophic).

Two families under Methanosarcinales represent aceticlastic (Methanosaetaceae) and hydrogenotrophic (Methanosarcinaceae) methanogens in the anaerobic digester. Their dominance is dependent on the concentrations of short chain fatty acids and ammonia on the sludge. Methanosarcinaceae are predominant when concentrations of short chain fatty acids and ammonia in

sludge are high. Conversely, Methanosaetaceae predominates when these concentrations are low [81].

Methanogens appear to use  $H_2$  more efficiently as shown by the low  $H_2$  concentration, if not absent, in biogas, as compared to substantial amount of acetate substrate in the sludge after biogasification. Accordingly, enhanced  $CH_4$  yield is observed in digesters added with hydrogen-producing bacterium *Caldicellulosyruptor saccharolyticus* [69]. The slow rate of acetate utilization compared with  $H_2$  utilization during methanogenesis can be attributed to the slower growth rate of aceticlastic methanogens than hydrogenotrophic methanogens rather than the capacity to utilize the substrate itself. Hence, a more efficient utilization of  $H_2$  than that of acetate may not be always true [62,69,78,82,83]. Biomass containing less protein that can be converted into ammonia during anaerobic digestion can be predominated by homoacetogens and hydrogenotrophic methanogens under thermophilic condition [81,84,85]. A possible shift to aceticlastic methanogens can be observed under mesophilic condition. Therefore, methanogen population dynamics can be influenced by several factors including substrate availability (acetate, short chain fatty acid, and ammonia) and operational conditions such as pH and temperature, among others [84].

In the sulfate-rich marine environment, sulfate-reducing bacteria (SRB) can outcompete hydrogenotrophic and aceticlastic methanogens because  $H_2S$  production is thermodynamically more favorable than  $CH_4$  using short chain fatty acids, and  $H_2/CO_2$ . Hence, only methylotrophic methanogens are commonly observed in the marine environment because of the inability of SRB to utilize methylated substrates [86,87]. Upon sulfate depletion, SRB forms a syntrophic relationship with hydrogenotrophic methanogens, making  $H_2$  as an electron sink and  $CH_4$  as an end product [88].

**Table 3**  
The elemental compositions of the selected and other genus of seaweeds.

Species	Elemental composition (% w/w dry)						Ref.
	C	H	O	N	S	C/N	
Selected Genus							
<i>C. racemosa</i>	38.03			2.06		18.45	[99]
				2.48		19.9	[100]
<i>C. scalpelliformis</i>	27.43			1.68		16.33	[99]
<i>C. veravelensis</i>	21.75			1.24		17.57	
<i>H. clathratus</i>					8.2		[101]
<i>T. conoides</i>						50.7	[102]
<i>T. ornate</i>				0.74		58.0	[100]
<i>S. polyceratum</i>	11.9			0.64		21.7	[103]
<i>S. patens</i>	40.18	5.22	33.85	2.00	0.98	20.09 <sup>a</sup>	[104]
<i>S. tenerrimum</i>	22.44	4.34		1.83	6.60	12.26 <sup>a</sup>	[105]
<i>U. capensis</i>				3.1			[106]
<i>U. fasciata</i>				1.44		25.7	[100]
<i>U. lactuca</i>				2.9			[106]
<i>U. prolifera</i>	29.7		50.1		2.97		[107]
<i>U. lactuca</i> (6 μM N)	37.0			1.66		26.02	[108]
<i>U. lactuca</i> (100 μM N)	38.13			4.09		10.88	
<i>U. rigida</i>				3.4			[106]
	28.7			1.3		22.1 <sup>a</sup>	[109]
<i>Ulva</i> spp. (High N)				3.24		8.72	[110]
<i>Ulva</i> spp. (Low N)				0.88		30.71	
Other Genus							
<i>Fucus serratus</i>	33.5	4.78	34.44	2.39	1.31	14.02 <sup>a</sup>	[111]
<i>F. vesiculosus</i>	32.88	4.77	35.63	2.53	2.44	13.00 <sup>a</sup>	
<i>Laminaria digitata</i>	31.59	4.85	34.16	0.90	2.44	36.53 <sup>a</sup>	
<i>L. hyperborea</i>	34.97	5.31	35.09	1.12	2.06	31.22 <sup>a</sup>	
<i>Macrocystis pyrifera</i>	27.3	4.08	34.8	2.03	1.89	13.45 <sup>a</sup>	

C – Carbon; H – Hydrogen; O – Oxygen; N – Nitrogen; S – Sulfur.

<sup>a</sup> Computed values by dividing N to C.

## 4. Biomethane potential of seaweed biomass

### 4.1. Properties of seaweed biomass

Most seaweeds are composed of phycocolloids. *Kappaphycus alvarezii* (Doty) Doty ex Silva has high  $\kappa$ -carrageenan polysaccharide [89], where as *Eucheuma denticulatum* (Burman) Collins & Hervey contains  $\iota$ -carrageenan [90]. *Sargassum* and *Padina* have 35% and 18.5% alginate, respectively [91]. Fucose and xylose are also present in *Sargassum* [92], whereas agar is the main component of *Gracilaria* [93]. In anaerobic digestion, the theoretical methane yield of a certain biomass can be obtained by determining elemental and proximate compositions. Different seaweeds contain different ratios of degradable components, which can significantly affect the efficiency of biogasification. Some important degradable components of the selected seaweeds *Sargassum* spp., *Turbinaria* spp., *Hydroclathrus* spp., *Caulerpa* spp., and *Ulva* spp. are discussed.

#### 4.1.1. Elemental components

The elements in organic matter are mainly carbon (C), hydrogen (H), oxygen (O), nitrogen (N), and sulfur (S). After moisture removal, the major compositions of biomass by mass are approximately 35–50% carbon and 40–45% oxygen [16]. The suggested optimum carbon to nitrogen ratio to achieve stable anaerobic digestion is between 25 and 30 [94]. A high amount of nitrogen is necessary for bacterial growth, especially during protein structure synthesis and nuclear matrix replication; however, an excessive amount may result in ammonia accumulation which has toxic

**Table 4**  
The proximate compositions of the selected genus of seaweeds.

Species	Proximate values (% w/w dry except moisture)						Ref.
	Lipids	CHO <sup>a</sup>	Protein	Fiber	Ash	Moisture	
Selected Genus							
<i>C. lentillifera</i>	0.86	59.27	12.49	3.17	24.21	25.31	[122]
	2.7	12.8	6.6		48.9		[123]
<i>C. racemosa</i>	19.1	83.2	18.3				[124]
	0.09	3.60	3.98	1.36	55.11		[125]
	3.8	16.6	6.8		42.2		[123]
	2.64	48.95	12.88		24.20	91.53	[99]
<i>C. scalpelliformis</i>	3.06	38.84	10.50		40.77	81.77	
<i>C. veravelensis</i>	2.80	37.23	7.77		33.70	87.88	
<i>H. clathratus</i>	2.18	82.26	6.39	2.70	6.47	59.63	[126]
	2.9	18.3	4.2		49.4		[123]
			8.6				[101]
<i>S. decurrens</i>	3.3	22.2	7.1		30.4		[123]
<i>S. filifolium</i>	4.0	21.4	10.2		28.2		
<i>S. horneri</i>	0.82	19.93	22.38		32.0	86.94	[127]
<i>S. longifolium</i>	8.2	16.8	18.65				[123]
<i>S. mangarevense</i>	3.4	24.9	13.2	17.9	60.6		[128]
<i>S. naozhouense</i>	1.06	47.73	11.20	4.83	35.18		[129]
<i>S. patens</i>					17.77	14.38	[104]
<i>S. tenerrimum</i>		8.20	0.86				[105]
	1.46	23.55	12.42				[130]
<i>S. wightii</i>	2.33	23.50	11–11.5				
<i>Sargassum</i> spp.	0.75	41.81	10.25	9.84	26.19	11.16	[58]
<i>T. conoides</i>	3.0	14.9	15.9				[131]
	1.9–2.1	23.9	11.5–12				[130]
	2.3	19.7	5.9		34.4		[123]
		31.83	8.6		34.30		[132]
<i>T. ornata</i>	2.2	24.6	9.2	13.5	39.8		[128]
<i>T. triquetra</i>	4.83	45.68	10.12		40.34	15.83	[133]
<i>U. lactuca</i>	1.6	23–24	3.25				[130]
	4.36	35.27	8.44				[134]
	1.64	14.6	7.06	55.4	21.3	10.6	[135]
<i>U. reticulata</i>	0.75	55.77	21.06	4.84	17.58	22.51	[122]
<i>Ulva</i> spp. (High N)		25.5	12.1	11.0			[110]
<i>Ulva</i> spp. (Low N)		51.4	3.6	4.3			

<sup>a</sup> CHO – Carbohydrates.



**Table 5**

The carbohydrate compositions (w/w dry) found on some of the selected genus of seaweeds.

Species	Compositions							Ref.
Brown seaweed	% Cel	% He	% Alg	% Man	% Fu	% La	% Lig	
<i>H. clathratus</i>			5.68	2.52		3.58		[136]
<i>S. filipendula</i>			14.4	0.8	26.0			[137]
<i>S. ilicifolium</i>				5				[138]
			30.80					[139]
<i>S. muticum</i>	2.2		16.9	7.7		0.3		[140]
<i>S. tenerrimum</i>				9.4				[138]
<i>S. wightii</i>			31.70	7.3				[141]
<i>Sargassum</i> spp.		25.73		5.04				[58]
<i>T. conoides</i>							19.27	[132]
				7.4				[142]
<i>T. ornata</i>			32.18	7.1				
<i>T. triquetra</i>			30.11					[133]
<i>T. turbinata</i>			24.6	0.2	32.1			[137]
Green seaweed	% Cel	% He	% Rh	% Xy	% Ua		% Lig	
<i>U. lactuca</i>	9.13	20.60					1.56	[143]
<i>U. lactuca</i> (Soluble fraction)			11.5	1.9	19			[114]

Cel – Cellulose; He – Hemicellulose; Alg – Alginic acid; Man – Mannitol; Fu – Fucoidan; La – Laminaran; Lig – Lignin; Rh – Rhamnose; Xy – Xylose; Ua – Uronic acid.

effects on methanogens [84,95,96]. Hence, the co-digestion of different biomasses are done to adjust the C/N ratio to an optimum value. The presence of a high amount of S as sulfate can also affect CH<sub>4</sub> production by encouraging the growth of SRB. These bacteria are more efficient in utilizing acetate and H<sub>2</sub>/CO<sub>2</sub> substrates than aceticlastic and hydrogenotrophic methanogens, respectively [78,97]. This may cause the production of high amount of H<sub>2</sub>S instead of CH<sub>4</sub>, consequently decreasing biogas quality and combustibility.

Table 3 shows the elemental composition of the different selected seaweeds. Methane potential can be computed by using the elemental composition of the biomass through the formula of Buswell and Mueller [98]:

$$C_nH_aO_b + \left(n - \frac{a}{4} - \frac{b}{2}\right)H_2O \rightarrow \left(\frac{n}{2} - \frac{a}{8} + \frac{b}{4}\right)CO_2 + \left(\frac{n}{2} + \frac{a}{8} - \frac{b}{4}\right)CH_4 \quad (1)$$

where *n*, *a*, and *b* are the mass fraction of elemental carbon, hydrogen and oxygen, respectively, on a dry weight basis of the biomass.

Computation of the theoretical methane yield using the elemental composition can overestimate the methane potential of a biomass because some amount of C, such as lignin, can be part of the non-degradable component of the biomass. Hence, proximate compositions are preferred for the methane potential computation of the biomass.

#### 4.1.2. Proximate compositions

Table 4 shows the proximate compositions of the selected seaweeds. Fresh seaweed is mainly composed of moisture and total solid (TS, dry matter). Moisture content varies among seaweeds, ranging from 10% to 90%. Volatile solid (VS), which is computed by subtracting ash from TS, primarily affects the methane potential of seaweeds. Among the VS components, the carbohydrates, proteins, and lipids of the biomass are used to compute for the theoretical methane yield. The constant values of carbohydrate (fiber and N-free extract), protein, and lipid as suggested by Karpenstein-Machan [112 in 113] are 395 l CH<sub>4</sub>/kg VS, 490 l CH<sub>4</sub>/kg VS, and 850 l CH<sub>4</sub>/kg VS, respectively.

Among the different carbohydrates components of the green seaweed *Ulva* (ulvan, α-cellulose, β-1,4-D-glucuronan and β-1,4-D-xyloglucan) [114–117], the sulfated ulvan can prevent the enzymatic digestion of α-cellulose [116,118,119], which may lead to incomplete hydrolysis. Brown seaweeds store carbohydrates as mannitol and laminarans, and their cell walls are composed of

alginates and fucoidans [120,121]. Although most of the VS of seaweeds are mainly composed of carbohydrates, the structures differ with that of terrestrial plants, which are mainly composed of cellulose and starch. Hence, the ability of the microorganisms that are conventionally used in biogas fermentation to anaerobically degrade seaweeds may be limited. Table 5 presents the different carbohydrates found in seaweeds.

#### 4.2. Methane fermentation of seaweed biomass

Many seaweed species have been studied as feedstock for methane fermentation. The theoretical methane yields of these seaweeds are high, but the actual methane yield of some species are low because of the presence of hydrolysis-resistant structural components or the absence of appropriate microorganisms that can breakdown the seaweed structures. Hence, the development of a suitable pretreatment for seaweed biomass is essential to maximize methane yield [144]. In the study of Bird et al. [2], *Sargassum fluitans* and *S. pteropleuron* yielded only 40% (450 l CH<sub>4</sub>/kg VS) and 35.7% (420 l CH<sub>4</sub>/kg VS) of their methane potential, respectively. Gupta et al. [145] discussed various approaches to enhance the methane yield of different biomass, such as co-digestion, as well as physical, chemical, and biological pretreatments. Physical pretreatment, specifically maceration, increases the methane yield of *Ulva lactuca* from 174 to 271 l CH<sub>4</sub>/kg VS, whereas thermal treatment at 130 °C increases methane yield to 187 l CH<sub>4</sub>/kg VS [146]. Without pretreatment, *Laminaria hyperborea* and *L. saccharina* produce 280 l CH<sub>4</sub>/kg VS and 230 l CH<sub>4</sub>/kg VS, respectively [147]. *Saccharina latissima* (*L. saccharina*) has low C/N ratio (8.8), but its methane yield (223 l CH<sub>4</sub>/kg VS) is enhanced by 20.18% and 16.59% with a 10-min steam explosion pretreatment at 130 °C and 160 °C, respectively [148]. Moreover, the co-digestion of *S. latissima* with steam-exploded wheat straw (210 °C, 10 min) to increase the C/N value to 30.2 improves its methane yield to 270 l CH<sub>4</sub>/kg VS [148].

The biomethane potential of *Ulva* sp. (196 l CH<sub>4</sub>/kg VS), *Gracilaria* sp. (182 l CH<sub>4</sub>/kg VS) and *Enteromorpha* sp. (154 l CH<sub>4</sub>/kg VS) decreases (167, 170 and 148 l CH<sub>4</sub>/kg VS, respectively) upon increasing TS input from 2.5% to 5% [149]. Furthermore, seaweed exhibit seasonal variation of methane yield. In *Laminaria digitata*, the highest actual methane yield (254.14 l CH<sub>4</sub>/kg VS) is obtained on July and the lowest (196.33 l CH<sub>4</sub>/kg VS) on March, corresponding to low and high ash content [150]. The seasonal variations in



the mannitol and laminaran content of *L. hyperborea* can also affect methane yield [151]. On the other hand, a two-day biological pretreatment of *L. japonica* (8.3 g Volatile Fatty Acid [VFA]/L), *Pachymeniopsis elliptica* (6.8 g VFA/L) and *Enteromorpha crinita* (4.4 g VFA/L) using *Vibrio harveyi* (15.6, 12.0 and 9.8 g VFA/L, respectively) and *V. alginolyticus* (~14, ~11.9 and ~7.5 g VFA/L, respectively) boosts VFA production, whereas chemical pretreatment using NaOH for 24 h is less effective (~12.8, ~9, and ~7.5 g VFA/L, respectively) [152].

The microflora of marine biomass can be used as microbial seed for anaerobic digestion [33], but operating conditions used in methane fermentation studies of seaweeds are patterned after biogas digesters running under terrestrial conditions (low salinity). Thus, the biogasification process utilizes washed biomass, freshwater liquid substrate, and conventional microbial inoculum. Operating conditions must be adapted to the requirements of marine microorganisms (thalassic), such as optimum salinity and temperature. With respect to anaerobic marine bacteria, *Clostridium halophilum* (degrades carbohydrates [78]), as well as *Halo-cella cellulolytica* [153 in 154] and *Sporohalobacter* [155 in 154] (both degrade cellulose), facilitates the anaerobic breakdown of cellulosic organic matter under high salinities. *Orenia marismortui*, *Halobacteroides halobius*, and *Sporohalobacter lortetii* metabolize glycogen and starch [156], and *Haloanaerobium praevalens* degrades pectin [154]. The presence of marine bacteria *Pseudomonas citrea*, which degrades alginate [157], and *Pseudomonas galatica* [158] and *Vibrio* species, which can both metabolize agar [159] and alginate [152], may increase the anaerobic degradation treatability of the seaweeds.

The innate ability of marine bacteria to effectively digest the unique structural components of seaweeds may result in higher methane yield if these organisms are used instead of the conventional inoculum. Hence, digester operating conditions should be adjusted to the needs of the marine bacteria. Furthermore, the use of marine bacteria as microbial seed may minimize the pretreatment needs for seaweed biomass because marine bacteria are generally more efficient in degrading seaweeds than the terrestrial counterpart that has been gradually developed for thalassic conditions. Preliminary results of studies on the anaerobic degradability of *Ulva* spp. show higher methane yield under thalassic conditions using marine microbial inoculum than that under freshwater conditions, wherein conventional microbial inoculum that is primarily obtained from the continuous fixed-bed biogas reactor is used [160]. Thus, the potential of household digester operation in coastal community, together with the accessibility of seaweed as biomass feedstock and seawater as the liquid substrate, can be demonstrated under thalassic conditions. Coastal communities can greatly benefit from these conditions especially in terms of convenience in handling and managing digesters. Although different pretreatments can greatly increase the anaerobic degradability of seaweed biomass, most chemical and physical pretreatments are not accessible for island folks. However, biological hydrolytic pretreatment using facultative marine microbial inoculum may be the most suitable and cheapest way for utilization by island communities.

## 5. Engaging coastal communities on green energy

Coastal communities are often among the poorest of the poor, with little to no access to basic social services. These conditions increase their vulnerability to the risks and negative impacts of climate change, adding more challenge to an already difficult life. Poverty alleviation requires multi-faceted effort, and must harmonize the complex issues of human well-being with environment and ecological sustainability, among others.

Sustainability in energy production and utilization by coastal populations can be achieved by using renewable energy sources. The major considerations include accessibility and cost in acquisition. The capacity and availability of seaweed biomasses as feedstock for energy production are demonstrated herein. This information can be used for institutionalizing green energy production for coastal populations.

Different biogas digesters with varying capacities have been installed in some regions of the Philippines. There were 45 digesters ( $\geq 500 \text{ m}^3$ : 1; 100–499  $\text{m}^3$ : 4; 10–29  $\text{m}^3$ : 24; 1–9  $\text{m}^3$ : 14; unknown: 2) in Region V; 25 digesters (all unknown capacities but were used for cooking/lighting) in Region IX; 13 digesters (30–99  $\text{m}^3$ : 2; 10–29  $\text{m}^3$ : 3; 1–9  $\text{m}^3$ : 8) in Region X; 48 digesters ( $\geq 500 \text{ m}^3$ : 1; 10–29  $\text{m}^3$ : 19; 1–9  $\text{m}^3$ : 21; unknown: 7) in Region XI; and 15 digesters (10–29  $\text{m}^3$ : 9; 1–9  $\text{m}^3$ : 6) in Region XIII [161,162]. However, the biogas technologies that are widely used today utilize freshwater which is usually scarce in coastal areas, especially in island communities, where rainwater is often the primary source of freshwater. Thus, the rationalization of freshwater use is a must. Given that biogas technology under saline conditions is feasible [33], the use of seawater, which is abundant in these communities, is more practical.

Although biogas technology in the Philippines is already developed, operational digesters are few and biogas capacities are low because of the limited accessibility of the communities to the technology [163] and expensive investment cost of digesters [164]. The appreciation of biogas technologies remains low in the country, especially at the grass roots level. The dissemination of information on biogas technology, specifically an easily understandable yet complete technical know-how, to the local government units and communities are lacking. A more directed, intentional, and localized approach such as information campaigns using local dialects may go a long way.

Operational digesters can be demonstrated in the community, with emphasis on its convenience in utilization and its long-term cheaper cost than that of conventional fossil fuels. Government incentives on using green energy may also be institutionalized to encourage further green energy production and utilization, especially biogas technology. Engaging the private sectors through public-private partnerships may also improve the accessibility and prolific adoption of the technology. To augment the operational cost and further increase profitability, the utilization of digester waste sludge as biofertilizer for mariculture or agriculture can also be studied. If a commercial biogas plant is preferred to small digesters, sustainable and managed harvesting of natural stocks and extensive mariculture of seaweeds for feedstock demand can be done by coastal folks as a source of additional income. The utilization of these seaweeds as feedstock for biogas production must also consider the potential demands on the high-priced natural products derived from the aforementioned seaweed species (e.g., alginate and fucoidan from the brown algae *Sargassum*, *Turbinaria*, and *Hydroclathrus*), such that the utilization of these resources is maximized and the goals of sustainable development are achieved. Although seaweed biomass resources are the focus of this review, the potential of dislodged sea grass biomasses that are commonly mixed with seaweed must be also considered.

## 6. Conclusions

The managed exploitation of underutilized renewable resources is necessary to encourage and maintain sustainable growth. At present, much of the seaweed resources remain underutilized in many coastal areas of the Philippines and elsewhere in the world despite its great potential to support sustainable development. These resources also hold great potential as alternative sources of biofuel, especially for the well-being of coastal

populations. The major advantages of seaweeds over its terrestrial counterpart are its shorter life cycle and easier, more cost-effective, and often environment-friendly cultivation. Five of the most viable Philippine seaweed species are the brown seaweeds *Sargassum* spp., *Turbinaria* spp., and *Hydroclathrus* spp., and green seaweeds *Caulerpa* spp. and *Ulva* spp. *Sargassum* spp. is the most abundant and well-studied among the species. The presence of extensive *Sargassum* beds throughout the country led to the development of an ecologically-sound cropping management for the natural stocks. *Turbinaria* spp. has the second largest natural bed in the country, but studies on its ecology and distribution are still limited. However, it is believed that *Turbinaria* may be sustainably harvested using the management scheme developed for *Sargassum*. Although seasonal, the extensive blooms of *Hydroclathrus* and *Ulva* species are correlated with eutrophication and thus can be utilized as biomass source, which also serves as a potential pollution remediation. At present, only *Caulerpa* has an established culture technology in the Philippines among the species presented herein. Culture technologies for the other seaweed species should be developed to sustainably produce biomass as feedstock for biogas production and avoid adverse effects on natural stocks (e.g., species extinction). Thus, research and development on this aspect is wanting.

Seaweed biomass is often treated as terrestrial feedstock. Seaweeds are washed to remove salts and conventional microbial inocula are used to initiate anaerobic digestion. Although the methane yield in this route showed promising results, the required pretreatment under similar digester conditions may pose a great challenge to island communities because the required resources are often limited and not easily accessible (e.g., freshwater). Hence, the utilization of biogas technology for these specific users may not be appealing. Thalassic biogas digesters that are adapted to island communities and use the aforementioned seaweeds may provide cheaper alternative and environment-friendly energy source. Aside from the benefits received by coastal populations from the energy supplied by this green technology, the society also benefits from the reduced carbon footprint because of non-reliance to fossil fuels, therefore helping in mitigating the negative impacts of climate change.

Among the suggested seaweeds, only *Sargassum* spp. and *Ulva* spp. have been extensively studied as resource for biomethane production. Research on the anaerobic digestion of *Turbinaria* spp., *Hydroclathrus* spp., and *Caulerpa* spp. are still needed. The optimum operating conditions that are suitable for the Philippine climate and appropriate biomass pretreatment that can be handled easily need immediate investigation. The utilization of post fermentation residue as biofertilizer for agriculture and mariculture, especially in seaweed farming, also requires further evaluation because nutrient enrichment may cause microalgal blooms. In addition, the benefits of using commercial or household biogas digesters should be locally demonstrated to encourage further the utilization of biogas technology. Support from the government is essential in bridging the technology to the communities.

## Acknowledgment

The corresponding author would like to acknowledge the support of the 2013 Research Grant from the Graduate School of Engineering, Nagoya University and the scholarship grant from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) in Japan. WJES and GCT Jr. would like to thank Mr. Christian Ace Guerta for helping with the literature search. The authors would like to acknowledge Ms. Najeeb Arabelle M. Rula for her assistance in the language check. This is the Marine Science

Institute (University of the Philippines) contribution no. 430 of WJES, GCT Jr., and MNEM.

## References

- [1] Campbell CJ. Oil crisis. United Kingdom: Multi-Science Publishing Co. Ltd.; 2005.
- [2] Bird KT, Chynoweth DP, Jerger DE. Effects of marine algal proximate composition on methane yields. *J Appl Ecol* 1990;2:207–313.
- [3] Chynoweth DP, Fannin KF, Srivastava VJ. Biological gasification of marine algae. In: Bird KT, Benson PH, editors. *Seaweed cultivation for renewable resources*. Amsterdam: Elsevier; 1987. p. 285–95.
- [4] Timilsina GR, Mevel S. Biofuels and climate change mitigation: a CGE analysis incorporating land-use change, World bank policy research working paper 5672. Washington, DC: The World Bank; 2011.
- [5] Rebhan E. Challenges for future energy usage. *Eur Phys J Spec Top* 2009;176: 53–80.
- [6] Sexton S, Rajagopal D, Zilberman D, Hochman G. Food versus fuel: how biofuels make food more costly and gasoline cheaper. *Agric Resour Econ Update* 2009;12:1–6.
- [7] Velimirov B, Field JG, Griffiths CL, Zoutendyk P. The ecology of kelp bed communities in the Benguela upwelling system, analysis of biomass and spatial distribution. *Helgol Wiss Meeresunter* 1977;30:495–518.
- [8] Gao Y, Skutsch M, Masera O, Pacheco P. A global analysis of deforestation due to biofuel development: a working paper 68. CIFOR, Bogor, Indonesia; 2011.
- [9] Harmeling S. Global climate risk index 2011. Germany: Germanwatch e.V., Germanwatch; 2010; 1–24.
- [10] Kreft S, Eckstein D. In: Kolboske B, Baum D, Kier G, editors. *Global climate risk index 2014*. Germany: Germanwatch e.V., Germanwatch; 2013. p. 1–28.
- [11] Corpuz PG. Philippine biofuels industry situation and outlook; 2012. Retrieved on July 2013 from: [http://gain.fas.usda.gov/Recent%20GAIN%20Publications/Biofuels%20Annual\\_Manila\\_Philippines\\_7-20-2012.pdf](http://gain.fas.usda.gov/Recent%20GAIN%20Publications/Biofuels%20Annual_Manila_Philippines_7-20-2012.pdf).
- [12] Energy Regulatory Commission-Philippines. ERC approves feed-in tariff rates. Retrieved on August 2013 from: <http://www.erc.gov.ph/pressrelease/ViewPressRelease/ERC-Approves-Feed-in-tariff-rates>.
- [13] Vasudevan P, Sharma S, Kumar A. Liquid fuel from biomass: an overview. *J Sci Ind Res India* 2005;64:822–31.
- [14] Padilla A. Pro-Arroyo landlords in congress to reap profits from biofuel act. *Bulatlat*, vol. VII, no. 3; February 18–24, 2007. Retrieved on July 2013 from: <http://www.bulatlat.com/news/7-3/7-3-biofuels.htm>.
- [15] Deublein D, Steinhauser A. *Biogas from waste and renewable sources: an introduction*. Weinheim: Wiley-VCH Verlag GmbH & Co. KGaA; 2008.
- [16] Chandra R, Takeuchi H, Hasegawa T. Methane production from lignocellulosic agricultural crop wastes: a review in context to second generation of biofuel production. *Renew Sustain Energy Rev* 2012;16:1462–76.
- [17] Central Intelligence Agency. *The World Factbook* 2013–14. Washington, DC; 2013. Retrieved on February 2014 from: <https://www.cia.gov/library/publications/the-world-factbook/fields/2060.html>.
- [18] Trono Jr. GC. Diversity of the seaweed flora of the Philippines and its utilization. *Hydrobiologia* 1999;398/399:1–6.
- [19] Trono Jr. GC. Field guide and atlas of the seaweed resources of the Philippines. Makati City, Philippines: Bookmark Inc.; 1997.
- [20] Guiry MD. In: Guiry MD, Guiry GM, editors. *AlgaeBase*, National University of Ireland. Galway: World-Wide Electronic Publication; 2013. <http://www.algaebase.org> (Retrieved on July 2013 from).
- [21] Montaña MNE, Rodriguez MRC, Balitaan RL. Ethnobotany of *Sargassum* spp. in the Philippines. *Coast Mar Sci* 2006;30:222–5.
- [22] Santiañez WJE, Trono Jr GC. Taxonomy of the Genus *Sargassum* (Fuciales, Phaeophyceae) from Alabat Island, Quezon, Northeastern Philippines. *Sci Diliman* 2013;25:29–50.
- [23] Ang PO. Phenology of *Sargassum siliquosum* J. Ag. and *S. paniculatum* J. Ag. (Sargassaceae, Phaeophyta) in the reef flat of Balibago (Calatagan, Philippines). In: Vivien MH, Salvat B, editors. *Proceedings of the fifth international coral reef congress*; 1985. 5. p. 51–7.
- [24] Ang PO. Phenology of *Sargassum* spp. in Tung Ping Chau Marine Park, Hong Kong SAR, China. *J Appl Phycol* 2006;18:629–36.
- [25] Largo DB, Ohno M. Phenology of two species of brown seaweeds, *Sargassum myriocystum* J. Agardh and *Sargassum siliquosum* J. Agardh (Sargassaceae, Fuciales) in Liloan, Cebu, in Central Philippines. *Bull Mar Sci Fish Kochi Univ* 1992;12:17–27.
- [26] Calumpung HP, Maypa AP, Magbanua M. Population and alginate quality assessment of four *Sargassum* species in Negros Island, Central Philippines. *Hydrobiologia* 1999;398/399:211–5.
- [27] Gillespie RD, Critchley AT. Phenology of *Sargassum* spp. (Sargassaceae, Phaeophyta) from Reunion Rocks, KwaZulu-Natal, South Africa. In: Kain JM, Brown MT, Lahaye M, editors. *Sixteenth international seaweed symposium*, *Hydrobiologia* 1999;398/399:201–10.
- [28] Arenas F, Fernandez C. Ecology of *Sargassum muticum* (Phaeophyta) on the North Coast of Spain: III. Reproductive ecology. *Bot Mar* 1998;41:209–21.
- [29] Atewerbehan M, Bruggemann JH, Breeman AM. Seasonal dynamics of *Sargassum ilicifolium* (Phaeophyta) on a shallow reef flat in the southern Red Sea (Eritrea). *Mar Ecol Prog Ser* 2005;292:159–71.

- [30] Rao AS, Rao MU. Seasonal growth pattern of *Sargassum polycystum* C. Agardh (Phaeophyta, Fucales) occurring at Visakhapatnam, east coast of India. *Indian J Mar Sci* 2002;31:26–32.
- [31] Tsuda R. Morphological, zonal and seasonal studies of two species of *Sargassum* on the reefs of Guam. *Proc Int Seaweed Symp* 1972;7:40–4.
- [32] Plouguerne E, Le Lann K, Connan S, Jechoux G, Deslandes E, Stiger-Pouvreau V. Spatial and seasonal variation in density, reproductive status, length and phenolic content of the invasive brown macroalga *Sargassum muticum* (Yendo) Fensholt along the coast of Western Brittany (France). *Aquat Bot* 2006;85:337–44.
- [33] Marquez GPB, Reichardt WT, Azanza RV, Klocke M, Montaño MNE. Thalassic biogas production from sea wrack biomass using different microbial seeds: cow manure, marine sediment and sea wrack-associated microflora. *Bioresour Technol* 2013;133:612–7.
- [34] Trono Jr GC, Tolentino GL. Studies on the management of *Sargassum* (Fucales, Phaeophyta) bed in Bolinao, Pangasinan, Philippines. *Korean J. Phycol.* 1993;8:249–57.
- [35] Ohno M, Ogawa H, Nakahara H, Orosco C. Ecological survey of *Sargassum* communities on the reefs of Central Visayas, Philippines. In: Umezaki I, editor. Scientific survey of marine algae and their resources in the Philippine islands, Technical report to the Ministry of Education, Science and Culture, Japan; 1987. p. 71–6.
- [36] Castro JRM. Taxonomy and distribution of the marine benthic algae of Taklong Island, Guimaras sub-province, Iloilo, Philippines. (Unpublished Master's thesis). Diliman: University of the Philippine; 1986.
- [37] Montes Jr HR. Temporal distribution and phenology of *Sargassum* population in Maydolong reef, Eastern Philippines. In: Calumpang HP, Menez EG, editors. Proceedings of the 2nd RP [Republic of the Philippines]-USA phycology symposium/workshop. Los Banos, Laguna (Philippines): PCAMRD; 1993. p. 127–44.
- [38] Ohno M, Noro T, Orosco CA. Ecological survey of *Sargassum* communities on the reefs of some Philippine Islands. In: Umezaki I, editor. Scientific Survey of Marine Algae and their resources in the Philippine Islands; 1989. p. 95–101.
- [39] Trono GC, Lluisma AO. Seasonality of standing crop of a *Sargassum* (Fucales, Phaeophyta) bed in Bolinao, Pangasinan, Philippines. *Hydrobiologia* 1990;204/205:331–8.
- [40] Corrales-Domingo AA. Taxonomy and distribution of the marine benthic algae of Ilocos Sur, Philippines. (Unpublished Master's thesis). Diliman: University of the Philippines; 1988.
- [41] Largo DB, Ohno M, Critchley A. Seasonal changes in the growth and reproduction of *Sargassum polycystum* C. Agardh and *Sargassum siliquosum* J. Agardh (Sargassaceae, Fucales) from Liloan, Cebu, in Central Philippines. *Jpn J Phycol* 1994;42:53–61.
- [42] Gaglac-Regis E. Distribution, growth and reproduction of two species of *Turbinaria* (Phaeophyta) in Pasacao, Camarines sur. (Unpublished Master's thesis). Diliman: University of the Philippines; 1982.
- [43] Ohno M, Largo DB, Trono Jr GC. A survey of standing crop, lengths of primary lateral branches and reproductive states of *Sargassum* communities on the reefs of the Philippine Islands. *Bull Mar Sci Fish Kochi Univ* 1995;15:67–78.
- [44] Ang Jr PO. Preliminary study on the sodium alginate contents of *Sargassum* spp. in Baliwago, Calatagan, Philippines. *Hydrobiologia* 1984;116/117:247–55.
- [45] Hurtado AQ, Ragaza AR. *Sargassum* studies in Currimaog, Ilocos Norte. I. Seasonal variation in the biomass of *Sargassum carpophyllum* J. Agardh, *Sargassum ilicifolium* (Turner) C. Agardh and *Sargassum siliquosum* J. Agardh (Phaeophyta, Sargassaceae). *Bot Mar* 1999;42:321–5.
- [46] Trono Jr GC. Seaweeds. In: Carpenter KE, Niem VH, editors. The living marine resources of the Western Central Pacific Volume 1: Seaweeds, corals, bivalves and gastropods. Rome: Food and Agriculture Organization of the United Nations; 1998. p. 19–99.
- [47] Ang Jr PO, Leung SM, Choi MM. A verification of reports of marine algae from the Philippines. *Philipp J Sci* 2013;142:5–49.
- [48] Velasquez GT. Studies on the marine algae of the Philippines. *Proc Eight Pac Sci Congr* 1957;4:556–62.
- [49] Gomez ED, Aliño PM, Yap HT, Licuanan WY. A review of the status of Philippine reefs. *Mar Pollut Bull* 1994;29:62–8.
- [50] San Diego-McGlone ML, Villanoy CL, Aliño PM. Nutrient mediated stress on the marine communities of a coastal lagoon (Puerto Galera, Philippines). *Mar Pol Bull* 1995;31(4–12):355–66.
- [51] Trono Jr. GC, Denila H. Studies on the pond culture of *Caulerpa*. 1987; *Phil Jour Sci, Monogr* 17:83–98.
- [52] Largo DB, Sembrano J, Hiraoka M, Ohno M. Taxonomic and ecological profile of 'green tide' species of *Ulva* (Ulvales, Chlorophyta) in central Philippines. *Hydrobiologia* 2004;512:247–53.
- [53] De Senerpont Domis LN, Fama P, Bartlett AJ, Prud'homme van Reine WF, Espinosa CA, Trono Jr GC. Defining taxon boundaries in members of the morphologically and genetically plastic genus *Caulerpa* (Caulerpaceae, Chlorophyta). *J Phycol* 2003;39:1019–37.
- [54] Trono Jr. GC. Field guide and atlas of the seaweed resources of the Philippines, 2. Makati City, Philippines: Bookmark Inc.; 2004.
- [55] Mata TM, Martins AA, Caetano NS. Microalgae for biodiesel production and other applications: a review. *Renew Sustain Energy Rev* 2010;14:217–32.
- [56] Brennan L, Owende P. Biofuels from microalgae – a review of technologies for production, processing, and extractions of biofuels and co-products. *Renew Sustain Energy Rev* 2010;14:557–77.
- [57] John RP, Anisha GS, Madhavan Nampoothiri K, Pandey A. Micro and macroalgal biomass: a renewable source for bioethanol. *Bioresour Technol* 2011;102:186–93.
- [58] Borines MG, de Leon RL, Cuello JL. Bioethanol production from the macro-algae *Sargassum* spp. *Bioresour Technol* 2013;138:22–9.
- [59] Hall DO. Biomass for energy in developing countries. *Geojournal* 1983;7: 5–14.
- [60] Stanley AM, Stanley DM, Dadu DW, Abah AM. Appraising the combustion of biogas for sustainable rural energy needs. *Afr J Environ Sci Technol* 2013;7: 350–7.
- [61] Chynoweth DP, Owens JM, Legrand R. Renewable methane from anaerobic digestion of biomass. *Renew Energy* 2001;22:1–8.
- [62] Nagamani B, Ramasamy K. Biogas production technology: an Indian perspective. *Curr Sci* 1999;77:44–55.
- [63] Nettmann E, Bergmann I, Pramschuer S, Mundt K, Plogsties V, Herrmann C, et al. Polyphasic analyses of methanogenic archaeal communities in agricultural biogas plant. *Appl Environ Microbiol* 2010;76:2540–8.
- [64] Yu Z, Schanbacher FL. Production of methane biogas as fuel through anaerobic digestion. In: Singh OV, Harvey SP, editors. Sustainable biotechnology. Netherlands: Springer; 2010. p. 105–27.
- [65] Demirel GN, Othman M. Two-phase thermophilic acidification and mesophilic methanogenesis anaerobic digestion of waste-activated sludge. *Environ Eng Sci* 2008;25:1291–300.
- [66] Elshahed MS. Microbiological aspects of biofuel production: current status and future directions. *J Adv Res* 2010;1:103–11.
- [67] Van de Werken HJG, Verhaart MRA, VanFossen AL, Willquist K, Lewis DL, Nichols JD, et al. Hydrogenomics of the extremely thermophilic bacterium *Caldicellulosiruptor saccharolyticus*. *Appl Environ Microbiol* 2008;74:6720–9.
- [68] Klass DL. Biomass for renewable energy, fuels, and chemicals. San Diego: Academic Press; 1998.
- [69] Bagi Z, Acs N, Bálint B, Horváth L, Dobó K, Perei KR, et al. Biotechnological intensification of biogas production. *Appl Microbiol Biotechnol* 2007;76: 473–82.
- [70] Shink B. Energetics of syntrophic cooperation in methanogenic degradation. *Microbiol Mol Biol Rev* 1997;61:262–80.
- [71] Boone DR, Bryant MP. Propionate-degrading bacterium, *Syntrophobacter wolnii* sp. nov. gen. nov., from methanogenic ecosystems. *Appl Environ Microbiol* 1980;40:626–32.
- [72] McInerney MJ, Bryant MP, Hespell RB, Costerton JW. *Syntrophomonas wolfei* gen. nov. sp. nov., an anaerobic, syntrophic, fatty acid-oxidizing bacterium. *Appl Environ Microbiol* 1981;41:1029–39.
- [73] Weiss A, Jérôme V, Burghardt D, Likke L, Peiffer S, Hofstetter EM, et al. Investigation of factors influencing biogas production in a large-scale thermophilic municipal biogas plant. *Appl Microbiol Biotechnol* 2009;84: 987–1001.
- [74] Winter JU, Wolfe RS. Methane formation from fructose by syntrophic associations of *Acetobacterium woodii* and different strains of methanogens. *Arch Microbiol* 1980;124:73–9.
- [75] Balk M, Weijma J, Stams AJM. *Thermotoga lettingae* sp. nov., a novel thermophilic, methanol-degrading bacterium isolated from a thermophilic anaerobic reactor. *Int J Syst Evol Microbiol* 2002;52:1361–8.
- [76] Kamagata Y, Mikami E. Diversity of acetotrophic methanogens in anaerobic digestion. In: Hattori T, Ishida Y, Maruyama Y, Morita RY, Uccida A, editors. Recent advances in microbial ecology. Tokyo, Japan: Japan Scientific Societies Press; 1989. p. 459–64.
- [77] Schnurer A, Schink B, Svensson BH. *Clostridium ultunense* sp. nov., a mesophilic bacterium oxidizing acetate in syntrophic association with a hydrogenotrophic methanogenic bacterium. *Int J Syst Bacteriol* 1996;46: 1145–52.
- [78] Lowe SE, Jain MK, Zeikus JG. Biology, ecology, and biotechnological applications of anaerobic bacteria adapted to environmental stresses in temperature, pH, salinity, or substrates. *Microbiol Rev* 1993;57:451–509.
- [79] Sakai S, Imachi H, Hanada S, Ohashi A, Harada H, Kamagata Y. *Methanocella paludicola* gen. nov., a methane-producing archaeon, the first isolate of the lineage 'Rice Cluster I', and proposal of the new archaeal order *Methanocellales* ord. nov. *Int J Syst Evol Microbiol* 2008;58:929–36.
- [80] Raskin L, Stromley JM, Rittmann BE, Stahl DA. Group-specific 16S rRNA hybridization probes to describe natural communities of methanogens. *Appl Environ Microbiol* 1994;60:1232–40.
- [81] Karakashev D, Batstone DJ, Angelidaki I. Influence of environmental conditions on methanogenic compositions in anaerobic biogas reactors. *Appl Environ Microbiol* 2005;71:331–8.
- [82] Angenent LT, Sung S, Raskin L. Methanogenic population dynamics during startup of a full-scale anaerobic sequencing batch reactor treating swine waste. *Water Res* 2002;36:4648–54.
- [83] Valentine DL. Biogeochemistry and microbial ecology of methane oxidation in anoxic environments: a review. *Antonie van Leeuwenhoek* 2002;81: 271–82.
- [84] Karakashev D, Batstone DJ, Trably E, Angelidaki I. Acetate oxidation is the dominant methanogenic pathway from acetate in the absence of *Methanosaetaceae*. *Appl Environ Microbiol* 2006;72:5138–41.
- [85] Krakat N, Westphal A, Schmidt S, Scherer P. Anaerobic digestion of renewable biomass: thermophilic temperature governs methanogen population dynamics. *Appl Environ Microbiol* 2010;76:1842–50.
- [86] Oremland RS, Marsh LM, Polcin S. Methane production and simultaneous sulphate reduction in anoxic, salt marsh sediments. *Nature* 1982;296:143–5.



- [87] Zinder SH. Ecology, physiology, biochemistry and genetics. In: Ferry JG, editor. Methanogens. USA: Springer; 1993. p. 128–206.
- [88] Plugge CM, Zhang W, Scholten JCM, Stams AJM. Metabolic flexibility of sulfate-reducing bacteria. *Front Microbiol* 2011;2:1–8.
- [89] Muñoz J, Freile-Pelegrin Y, Robledo D. Mariculture of *Kappaphycus alvarezii* (Rhodophyta, Solieriaceae) color strains in tropical waters of Yucatan, Mexico. *Aquaculture* 2004;239:161–77.
- [90] Funami T, Hiroe M, Noda S, Asai I, Ikeda S, Nishinari K. Influence of molecular structure imaged with atomic force microscopy on the rheological behavior of carrageenan aqueous systems in the presence or absence of cations. *Food Hydrocolloids* 2007;21:617–29.
- [91] Omar S, Ahmad N, Ahmad F. Composition of alginates from brown seaweeds, *Sargassum* and *Padina* spp. *Pertanika* 1988;11:79–85.
- [92] Marín A, Casas-Valdez M, Carrillo S, Hernández H, Monroy A, Sanginés L, et al. The marine algae *Sargassum* spp. (Sargassaceae) as feed for sheep in tropical and subtropical regions. *Rev Biol Trop* 2009;57:1271–81.
- [93] Givernaud T, El Gourji A, Mouradi-Givernaud A, Lemoine Y, Chiadmi N. Seasonal variations of growth and agar composition of *Gracilaria multipartite* harvested along the Atlantic coast of Morocco. *Hydrobiologia* 1999;398/399:167–72.
- [94] Mital KM. Biogas systems: principles and applications. New Delhi: New Age International (P) Limited Publishers; 1996; 412.
- [95] Sossa K, Alarcon M, Aspe E, Urrutia H. Effect of ammonia on the methanogenic activity of methylaminotrophic methane producing Archaea enriched biofilm. *Anaerobe* 2004;10:13–8.
- [96] Zhou H, Qiu G. Inhibitory effect of ammonia nitrogen on specific methanogenic activity of anaerobic granular sludge. *J Cent South Univ Technol* 2006;13:63–7.
- [97] King GM. Methanogenesis from methylated amines in a hypersaline algal mat. *Appl Environ Microbiol* 1988;54:130–6.
- [98] Buswell AM, Mueller HF. Mechanism of methane fermentation. *Ind Eng Chem* 1952;44:550–2.
- [99] Kumar M, Gupta V, Kumari P, Reddy CRK, Jha B. Assessment of nutrient composition and antioxidant potential of Caulerpaceae seaweeds. *J Food Compos Anal* 2011;24:270–8.
- [100] Lin H-J, Wu C-Y, Kao S-J, Kao W-Y, Meng P-J. Mapping anthropogenic nitrogen through point sources in coral reefs using  $\delta^{15}\text{N}$  in macroalgae. *Mar Ecol Prog Ser* 2007;335:95–109.
- [101] Wang H, Chiu LCM, Ooi VEC, Ang Jr PO. A potent antitumor polysaccharide from the edible brown seaweed *Hydroclathrus clathratus*. *Bot Mar* 2010;53:265–74.
- [102] Mayakun J, Kim JH, Lapointe BE, Prathep A. Algal response to nutrient enrichment; insights into growth and chemical concentrations in two algae, *Halimeda macroloba* Decaisne and *Turbinaria conoides* (J. Agardh) Kützting. *Songklanakarin J Sci Technol* 2013;35:389–95.
- [103] Lapointe BE. Nutrient thresholds for bottom-up control of macroalgal blooms on coral reefs in Jamaica and southeast Florida. *Limnol Oceanogr* 1997;42:1119–31.
- [104] Li D, Chen L, Xu D, Zhang X, Ye N, Chen F, et al. Preparation and characteristics of bio-oil from the marine brown alga *Sargassum patens* C. Agardh. *Bioresour Technol* 2012;104:737–42.
- [105] Vijayabaskar P, Vaseela N. In vitro antioxidant properties of sulfated polysaccharide from brown marine algae *Sargassum tenerrimum*. *Asian Pac J Trop Dis* 2012;S890–6.
- [106] Shuuluka D, Bolton JJ, Anderson RJ. Protein content, amino acid composition and nitrogen-to-protein conversion factors of *Ulva rigida* and *Ulva capsensis* from natural populations and *Ulva lactuca* from an aquaculture system, in South Africa. *J Appl Phycol* 2013;25:677–85.
- [107] Michalak I, Chojnacka K. Edible macroalgae *Ulva prolifera* as microelemental feed supplement for livestock: the fundamental assumptions of the production method. *World J Microbiol Biotechnol* 2009;25:997–1005.
- [108] Nielsen MM, Bruhn A, Rasmussen MB, Olesen B, Larsen MM, Møller HB. Cultivation of *Ulva lactuca* with manure for simultaneous bioremediation and biomass production. *J Appl Phycol* 2012;24:449–58.
- [109] Pinchetti JLG, Fernández EDC, Díez PM, Reina GG. Nitrogen availability influences the biochemical composition and photosynthesis of tank-cultivated *Ulva rigida* (Chlorophyta). *J Appl Phycol* 1998;10:383–9.
- [110] Habig C, DeBusk TA, Ryther JH. The effect of nitrogen content on methane production by the marine algae *Gracilaria tikvahiae* and *Ulva* sp. *Biomass* 1984;4:239–51.
- [111] Ross AB, Jones JM, Kubacki ML, Bridgeman T. Classification of macroalgae as fuel and its thermochemical behavior. *Bioresour Technol* 2008;99:6494–504.
- [112] Karpenstein-Machan M. Energiepflanzenbau für Biogasanlagenbetreiber (Energy crop for biogas plant operators). Frankfurt am Main, Germany: DLG-Verlags-GmbH; 2005.
- [113] Amon T, Amon B, Kryvoruchko V, Machmuller A, Hopfner-Sixt K, Bodiroza V, et al. Methane production through anaerobic digestion of various energy crops grown in sustainable crop rotations. *Bioresour Technol* 2007;98:3204–12.
- [114] Lahaye M, Jegou D. Chemical and physical-chemical characteristics of dietary fibers from *Ulva lactuca* (L.) Thuret and *Enteromorpha compressa* (L.) Grev. *J Appl Phycol* 1993;5:195–200.
- [115] Lahaye M, Ray B. Cell-wall polysaccharides from the marine green alga *Ulva "rigida"* (Ulvales, Chlorophyta) – NMR analysis of ulvan oligosaccharides. *Carbohydr Res* 1996;283:161–73.
- [116] Lahaye M, Brunel M, Bonnin E. Fine chemical structure analysis of oligosaccharides produced by an ulvan-lyase degradation of the water-soluble cell-wall polysaccharides from *Ulva* sp. (Ulvales, Chlorophyta). *Carbohydr Res* 1997;304:325–33.
- [117] Yamamoto M. Physicochemical studies on sulfated polysaccharides extracted from seaweeds at various temperatures. *Agric Biol Chem* 1980;44:589–93.
- [118] Lahaye M. NMR spectroscopic characterization of oligosaccharides from two *Ulva rigida* ulvan samples (Ulvales, Chlorophyta) degraded by lyase. *Carbohydr Res* 1998;314:1–12.
- [119] Alves A, Sousa RA, Reis RL. A practical perspective on ulvan extracted from green algae. *J Appl Phycol* 2013;25:407–24.
- [120] Evans LV, Simpson M, Callow ME. Sulphated polysaccharide synthesis in brown algae. *Planta* 1973;110:237–52.
- [121] Usov AI, Smirnova GP, Klochova NG. Polysaccharides of algae: 55.<sup>1</sup> Polysaccharide composition of several brown algae from Kamchatka. *Russ J Bioorg Chem* 2001;27:395–9.
- [122] Ratana-arporn P, Chirapart A. Nutritional evaluation of tropical green seaweeds *Caulerpa lentillifera* and *Ulva reticulata*. *Kasetsart J (Nat Sci)* 2006;40:75–83.
- [123] Renaud SM, Luong-Van JT. Seasonal variation in the chemical composition of tropical Australian marine macroalgae. In: Anderson R, Brodie J, Onsenoyen E, Critchley AT, editors. Eighteenth international seaweed symposium. Netherlands: Springer; 2007. p. 155–61.
- [124] Rameshkumar S, Ramakritinan CM, Eswaran K, Yokeshbabu M. Proximate composition of some selected seaweeds from Palk bay and Gulf of Mannar, Tamilnadu, India. *Asian J Biomed Pharm Sci* 2012;3:1–5.
- [125] Robledo D, Freile Pelegrin Y. Chemical and mineral composition of six potential edible seaweed species of Yucatán. *Bot Mar* 1997;40:301–6.
- [126] Zafar M, Chowdhury SMRA. Water quality and biochemical components of *Hydroclathrus clathratus* in the Tidal shore area of St. Martin's Island, Bangladesh. *Int J Phycol Phycochem* 2009;5:7–10.
- [127] Hossain Z, Kurihara H, Takahashi K. Biochemical composition and lipid compositional properties of the brown alga *Sargassum horneri*. *Pak J Biol Sci* 2003;6:1497–500.
- [128] Zubia M, Payri CE, Deslandes E, Guezennec J. Chemical composition of attached and drift specimens of *Sargassum mangroveense* and *Turbinaria ornata* (Phaeophyta: Fuciales) from Tahiti, French Polynesia. *Bot Mar* 2003;46:562–71.
- [129] Peng Y, Xie E, Zheng K, Fredimoses M, Yang X, Zhou X, et al. Nutritional and chemical composition and antiviral activity of cultivated seaweed *Sargassum naozhouense* Tseng et Lu. *Mar Drugs* 2013;11:20–32.
- [130] Manivannan K, Thirumaran G, Karthikai Devi G, Hemalatha A, Anantharaman P. Biochemical composition of seaweeds from Mandapam Coastal Regions along Southeast Coast of India. *Am-Eurasian J Bot* 2008;1:32–7.
- [131] Narasimman S, Murugaiyan K. Proximate composition of certain selected marine macro-algae from Mandapam Coastal Region (Gulf of Mannar), Southeast Coast of Tamil Nadu. *Int J Pharm Biol Arch* 2012;3:918–21.
- [132] Shyamali S, De Silva M, Savitri Kumar N. Carbohydrate constituents of marine algae of Sri Lanka. Part III. Composition of the carbohydrates extracted from the brown seaweed *Turbinaria conoides*. *J Natl Sci Counc Sri Lanka* 1988;16:201–8.
- [133] Omar HH, Abdullatif BM, El-Kazan MM, El-Gendy AM. Red sea water and biochemical composition of seaweeds at Southern Coast of Jeddah, Saudi Arabia. *Life Sci J* 2013;10:1073–80.
- [134] Chakraborty S, Santra SC. Biochemical composition of eight benthic algae collected from Sunderban. *Indian J Mar Sci* 2008;37:329–32.
- [135] Wong KH, Cheung PCK. Nutritional evaluation of some subtropical red and green seaweeds Part I – proximate composition, amino acid profiles and some physico-chemical properties. *Food Chem* 2000;71:475–82.
- [136] Awad NE, Motawe HM, Selim MA, Matloub AA. Antitumorigenic polysaccharides isolated from the brown algae, *Padina pavonia* (L.) Gaill. and *Hydroclathrus clathratus* (C. Agardh) Howe. *Med Aromat Plant Sci Biotechnol* 2009;3:6–11.
- [137] García-Ríos V, Ríos-Leal E, Robledo D, Freile-Pelegrin Y. Polysaccharides composition from tropical brown seaweeds. *Phycol Res* 2012;60:305–15.
- [138] Varier NS, Sadasivan Pillai K. Mannitol from *Sargassum* seaweeds. II. Optimum conditions for extraction of alginic acid from *Sargassum* seaweeds of Cape Gomorin. *Bull Cent Res Inst* 1952;2:39.
- [139] Chennubhotla VSK, Kaliaperumal N, Kalimuthu S, Selvaraj M, Ramalingam JR, Najnuddin M. Seasonal changes in growth and alginic acid and mannitol contents in *Sargassum ilicifolium* (Turner) J. Agardh and *S. myriocystum* J. Agardh. *Indian J Mar Sci* 1982;11:195–6.
- [140] Gorham J, Lewey SA. Seasonal changes in the chemical composition of *Sargassum muticum*. *Mar Biol* 1984;80:103–7.
- [141] Umamaheswara Rao M. Seasonal variations in growth, alginic acid and mannitol contents of *Sargassum wightii* and *Turbinaria conoides* from the Gulf of Mannar, India. In: Proceedings of the 6th International Seaweed Symposium; 1969. p. 579–84.
- [142] Umamaheswara Rao M, Kalimuthu S. Changes in mannitol and alginic acid contents of *Turbinaria ornata* (Turner) J. Agardh in relation to growth and fruiting. *Bot Mar* 1972;15:57–9.
- [143] Yaich H, Garna H, Besbes S, Paquot M, Blecker C, Attia H. Chemical composition and functional properties of *Ulva lactuca* seaweed collected in Tunisia. *Food Chem* 2011;128:895–901.
- [144] Dębowski M, Zieliński M, Grola A, Dudek M. Algae biomass as an alternative substrate in biogas production technologies – review. *Renew Sustain Energ Rev* 2013;27:596–604.

- [145] Gupta P, Singh RS, Sachan A, Vidyarthi AS, Gupta A. A re-appraisal on intensification of biogas production. *Renew Sustain Energy Rev* 2012;16:4908–16.
- [146] Bruhn A, Dahl J, Nielsen HB, Nikolaisen L, Rasmussen MB, Markager S, et al. Bioenergy potential of *Ulva lactuca*: biomass yield, methane production and combustion. *Bioresour Technol* 2011;102:2595–604.
- [147] Hanssen JF, Indergaard M, Østgaard K, Bævre OA, Pedersen TA, Jensen A. Anaerobic digestion of *Laminaria* spp. and *Ascophyllum nodosum* and application of end products. *Biomass* 1987;14:1–13.
- [148] Vivekanand V, Eijssink VGH, Horn SJ. Biogas production from the brown seaweed *Saccharina latissima*: thermal pretreatment and codigestion with wheat straw. *J Appl Phycol* 2012;24:1295–301.
- [149] Gonçalves PR, Costa JC, Nobre A, Alves MM. Biomethanation potential of macroalgae *Ulva* spp. and *Gracilaria* spp. and in co-digestion with wasted activated sludge. *Bioresour Technol* 2012;114:320–6.
- [150] Adams JMM, Toop TA, Donnison IS, Gallagher JA. Seasonal variation in *Laminaria digitata* and its impact on biochemical conversion routes to biofuels. *Bioresour Technol* 2011;102:9976–84.
- [151] Horn SJ, Østgaard K. Alginate lyase activity and acidogenesis during fermentation of *Laminaria hyperborea*. *J Appl Phycol* 2001;13:143–52.
- [152] Pham TN, Um Y, Yoon HH. Pretreatment of macroalgae for volatile fatty acid production. *Bioresour Technol* 2013;146:754–7.
- [153] Zhilina TN, Zavarzin GA, Bulygina ES, Kevbrin VV, Osipov GA, Chumakov KM. Ecology, physiology and taxonomy studies on a new taxon of Haloanaerobiaceae, *Haloancola saccharolytica* gen. nov., sp. nov. *Syst Appl Microbiol* 1992;15:275–84.
- [154] Ollivier B, Caumette P, Garcia J-L, Mah RA. Anaerobic bacteria from hypersaline environment. *Microbiol Rev* 1994;58:27–38.
- [155] Mathrani IM, Ollivier BM, Boone DR, Mah RA. Enrichment and enumeration of methanogenic, sulfate-reducing, and cellulolytic bacteria from lake Retba, a hypersaline lake in Senegal, West Africa. In: Abstract in 87th Annual Meeting of American Society of Microbiology. Washington, DC: American Society for Microbiology; 1987; 184.
- [156] Moune S, Eatock C, Matheron R, Willison JC, Hirschler A, Herbert R, et al. *Orenia salinaria* sp. nov., a fermentative bacterium isolated from anaerobic sediments of Mediterranean salterns. *Int J Syst Evol Microbiol* 2000;50:721–9.
- [157] Alekseeva SA, Bakunina IY, Nedashkovskaya OI, Isakov II, Mikhailov VV, Zvyagintseva TN. Intracellular alginolytic enzymes of the marine bacterium *Pseudoalteromonas citrea* KMM3297. *Biochemistry* 2004;69:262–9.
- [158] Zhang C, Kim S-K. Research and application of marine microbial enzymes: status and prospects. *Mar Drugs* 2010;8:1920–34.
- [159] Sugano Y, Matsumoto T, Kodama H, Noma M. Cloning and sequencing of agaA, a unique agarase 0107 gene from a marine bacterium, *Vibrio* sp. strain JT0107. *Appl Environ Microbiol* 1993;59:3750–6.
- [160] G.P.B. Marquez, H. Takeuchi and T. Hasegawa, Biogas production of biologically and chemically-pretreated seaweed, *Ulva* spp., under different conditions: freshwater and thalassic. *Bioresour Technol*, in preparation.
- [161] Department of Science and Technology (DOST), Republic of the Philippines, 2008. ([www.dost.gov.ph](http://www.dost.gov.ph)).
- [162] International Institute for Energy Conservation (IIEC), Eastern Research Group, Inc., PA Consulting Group. Resource assessment for livestock and agro-industrial wastes-Philippines; 2009. Retrieved on February 2014 from: ([https://www.globalmethane.org/documents/ag\\_philippines\\_res\\_assessment.pdf](https://www.globalmethane.org/documents/ag_philippines_res_assessment.pdf)).
- [163] The Japan Institute of Energy. The Asian biomass handbook, A guide for biomass production and utilization. In: Yokoyama S, Matsumura Y, Ando S, Sakanishi K, Sano H, Minowa T, et al., editors. Support project for building Asian-partnership for environmentally conscious agriculture; 2008.
- [164] Teune B, Orprecio J, Dalusung A, Yeneza G. Feasibility study of a national biogas programme on domestic biogas in the Philippines. SNV Netherlands Development Organisation and Winrock International; April 2010. Retrieved on July 2013 from: ([http://www.snvworld.org/sites/www.snvworld.org/files/publications/feasibility\\_study\\_of\\_a\\_national\\_domestic\\_biogas\\_programme\\_the\\_philippines\\_2010.pdf](http://www.snvworld.org/sites/www.snvworld.org/files/publications/feasibility_study_of_a_national_domestic_biogas_programme_the_philippines_2010.pdf)).